

1. Reflection about line $x = 0$ is **horizontal**, so replace $x \rightarrow -x$. Then, for horizontal translation, replace $x \rightarrow x + 3$, giving us $y = f[-(x + 3)]$ which simplifies to $y = f(-x - 3)$.

ANSWER: **D**

- NR #1 For $g(x)$, the transformation is to take all points on $f(x)$ and

- Horizontally shift 2 left So subtract 2 from all x -coords
- Vertically shift k up So add "k" to all y -coords

The DOMAIN of $f(x) = \sqrt{x+3} - 1$ is...

Can't sq root negatives $x + 3 \geq 0$
 $x \geq -3$ $\xrightarrow{\text{shift 2 left}}$ $x \geq -5$

The RANGE of $f(x) = \sqrt{x+3} - 1$ is...

Basic graph of $y = \sqrt{x}$ shifted 1 down $y \geq -1$ $\xrightarrow{\text{shift "k" up}}$ $y \geq -1 + k$
 Simplifies to: $y \geq k - 1$

ANSWER: **94**

- NR #2 Apply vert. str. factor of 3 to $2x^2 + 1$, to get $y = 6x^2 + 3 + 2$

$y = 3(2x^2 + 1)$
 $y = 6x^2 + 3$


ANSWER: **65**

- NR #3 Jagmeet sketches the graph of $h(x) = 4f(-x) - 3$

All pts $(x, y) \rightarrow (-x, 4y - 3)$

First (farthest left) pt on graph of $f(x)$ $(-4, 2) \rightarrow (4, 5)$
 $(0, 3) \rightarrow (0, 9)$
 $(3, -1) \rightarrow (-3, -7)$
 $(5, 0) \rightarrow (-5, -3)$

Plot each of these points!



So, domain is $[-5, 4]$ and range is $[-7, 9]$
 convert to CODES $\downarrow \downarrow \downarrow \downarrow$
 $3 \quad 4 \quad 2 \quad 9$

ANSWER: **3429**

5. On $f(x)$ the x -coord of P is 3
 Horiz. str. by a factor of 3 changes this to 9
 Then horiz. shift 9 units right changes this to $m = 18$

ANSWER: **D**

7. The domain of $f(x)$ is $x \geq 9$
 Think: Basic graph of $y = \sqrt{x}$ (domain $x \geq 0$) shifted 9 units right

Then we apply the mapping rule
 $3(9) - 6$ to get $x \geq 21$

ANSWER: **C**

8. The total width of $f(x)$ is 6 units. ($x = 2$ to $x = 8$)
 The total width of $g(x)$ is 1.5 units. ($x = 0.5$ to $x = 2$)
 \rightarrow Therefore $g(x)$ is $1/4^{\text{th}}$ as wide ($1.5 \div 6$)

HORIZ STR. FACTOR OF $1/4$ b value in equation is 4 (reciprocal)

2. Transformations for $g(x)$ are a vertical reflection (replace $y \rightarrow -y$) a vertical stretch, and a vertical shift one unit up.

So all pts $(x, y) \rightarrow (x, -2y + 1)$

Now, the RANGE of $f(x)$ (shown) is $[-4, \infty)$

Which on $g(x)$ becomes $(-\infty, 9]$

ANSWER: **A**

$(-2)(-4) + 1$

3. Transformations for $h(x)$ are a horizontal stretch, factor of 2, and a horiz. translation 4 units right.

*remember to first FACTOR: $h(x) = f[\frac{1}{2}(x - 4)]$

Now, the DOMAIN of $f(x)$ (shown) is $[-5, \infty)$

Which on $g(x)$ becomes $[6, \infty)$

ANSWER: **C**

$(2)(-5) + 4$

4. The largest x -intercept on $h(x)$ is -1 . After a horiz. str. factor of 2 and shift 4 right, that becomes

$x = 2$
 $(2)(-1) + 4$

ANSWER: **D**

6. An equation for $k(x)$ can be found by replacing, in the equation for $f(x)$

then, in the resulting expression $\frac{1}{3}x$... Replacing x with $x - 9$
 $\frac{1}{3}x \rightarrow \frac{1}{3}(x - 9)$
 Horiz str. factor of 3 Horiz shift 9 right

So net effect: $k(x) = f\left[\frac{1}{3}(x - 9)\right]$

ANSWER: **B**

The total height of $f(x)$ is 3 units. ($y = 0$ to $y = 3$)

The total height of $g(x)$ is 9 units. ($y = 2$ to $y = 11$)

\rightarrow Therefore $g(x)$ is 3 times "taller" VERT STR. FACTOR OF 3 BUT there is also a vertical shift, k .

Think: $g(x)$ "should" have invariant points on the x -axis, from the vertical stretch. (at the MIN)

However the MIN y -value of $g(x)$ is 2

ANSWER: **A**

\rightarrow Therefore k is 2 units up

- 9.** Reflection about line $x = 0$ replace $x \rightarrow -x$. **AND** Reflection about line $y = 0$ / think of one of two ways. Either replace $y \rightarrow -x$ or *make the entire expression negative*.

$$g(x) = -[2(-x)^2 - 3(-x) + 5]$$

For Vert. Refl. For Horiz. Refl. ($-$ sign inside)
(include square brackets around all terms)

$$g(x) = -[2x^2 + 3x + 5]$$

$$g(x) = -2x^2 - 3x - 5$$

ANSWER: **A**

- NR #5** Many options! Use the "furthest left" point on each graph.

On $f(x)$ the furthest left point is **6 horiz. units** from the y -axis, on $g(x)$ it's **9 horiz. units**.



→ Horiz. str, factor of $9/6$
reduces to $\frac{3}{2}$

On $f(x)$ the highest point is **2 vert. units** from the x -axis, on $g(x)$ it's **1 unit**.

→ vert. str, factor of $\frac{1}{2}$

SO... $a = \frac{1}{2} \leftarrow m$
 $\frac{3}{2} \leftarrow n$
(vert. str.)

AND... $b = \frac{2}{3} \leftarrow p$
 $\frac{1}{2} \leftarrow q$
(reciprocal of horiz. str)



ANSWER: **1223**

- NR #4** Start with $f(x)$ **DOMAIN** is $[-2, \infty)$ and **RANGE** is $[-4, \infty)$
"left to right" "bottom to top"

∗ $g(x)$ involves a Vert. Str, factor of $3/2$, a horiz. reflection, and a shift 2 right and 2 up.

All pts $(x, y) \rightarrow (-x - 2, \frac{3}{2}y + 2)$
affects domain affects range

DOMAIN of $g(x)$ $-(-2) - 2 \Rightarrow x \leq 0$ REF. #1
(reverse direction / graph points LEFT now)

RANGE of $g(x)$ $\frac{3}{2}(-4) + 2 \Rightarrow y \geq -4$ REF. #7

∗ $h(x)$ is the INVERSE, $(x, y) \rightarrow (y, x)$ Domain and range switch

DOMAIN of $h(x) \Rightarrow x \geq -4$ REF. #5
the RANGE of $f(x)$

RANGE of $h(x) \Rightarrow y \geq -2$ REF. #8
the DOMAIN of $f(x)$


ANSWER: **1758**

- 10.** First apply vertical reflection
 $= -[2x^2 + 3x - 5]$ - in front of entire expression
 $= -2x^2 - 3x + 5$


Then apply horiz shift 1 left Replace x with " $x + 1$ "
 $= -2(x + 1)^2 - 3(x + 1) + 5$
 $= -2(x^2 + 2x + 1) - 3x - 3 + 5$
 $= -2x^2 - 4x - 2 - 3x + 2$
 $= -2x^2 - 7x$

ANSWER: **B**

- NR #6** First determine vertical stretch (which is " a ") and horiz. stretch (which is reciprocal of " b ")



$f(x)$ is 8 units "wide"
(left to right ... $x = 4$ to $x = 12$)



and $g(x)$ is 6 units "wide"
(left to right ... $x = 3$ to $x = 9$)


and $g(x)$ is 21 units "tall"

→ Vert. str, factor of $21/15$
Decimal form is **1.4**



ANSWER: **1408**

→ Horiz. str, factor of $6/8$ Decimal form rounds to **0.8**

- NR #7** The graph of $g(x)$ can be obtained by switching all coordinates $(x, y) \rightarrow (y, x)$



- 1) There are TWO invariant points 2) The # of y-int on $g(x)$ is equal to the # of x-ints on $f(x) \rightarrow 3$



ANSWER: **233**

3) ... and the largest x -coord is **3** (the largest y -coord on $f(x)$)

11. Switch x and y , then isolate y

first re-write $f(x)$ in terms of y

$$y = \sqrt{x+4} - 1$$

$$x = \sqrt{y+4} - 1 \Rightarrow x+1 = \sqrt{y+4} \Rightarrow (x+1)^2 = (\sqrt{y+4})^2$$

$$\Rightarrow (x+1)^2 = y+4 \Rightarrow y = (x+1)^2 - 4 \quad \text{ANSWER: B}$$

12. The domain of $g(x)$ is the range of $f(x)$

Range of $f(x)$ is $y \geq -1$


So domain of $g(x)$ is $x \geq -1$

Graph is a "half-parabola"

And the y -int of $g(x)$ is $y = -3$

(The x -int of $f(x)$)

ANSWER: A



WR Question 1


(a) Vert. str. by factor of 3, then vert. translation 4 units right

$$(x, y) \rightarrow (x+4, 3y)$$

$$(-4, 3) \rightarrow (0, 9)$$

$$(-2, 1) \rightarrow (2, 3)$$

$$(0, 3) \rightarrow (4, 9)$$



(b) Equation of $f(x)$ $y = a(x+2)^2 + 1$ $\Rightarrow (3) = a((0)+2)^2 + 1$

$$\Rightarrow 2 = a(2)^2$$

$$\Rightarrow a = \frac{2}{4} \Rightarrow a = \frac{1}{2}$$

So, equation of $f(x)$ is

$$f(x) = \frac{1}{2}(x+2)^2 + 1$$

Use any other pt on the graph to solve for a (Such as $(0, 3)$!)

For equation of $g(x)$ we could use a similar method (sub in vertex then solve for "a")

OR we could simply apply the transformations to the equation of $f(x)$

$$g(x) = 3 \left[\frac{1}{2}((x-4)+2)^2 + 1 \right]$$

horiz. translation 4 right

Vert. str. factor of 3

So, equation of $g(x)$ is

$$g(x) = \frac{3}{2}(x-2)^2 + 3$$

(c) Here we saw that horizontally translating the vertex 4 right achieved the same result as horizontally reflecting the graph of $y = f(x)$.

This can be verified by applying the horiz. reflection to the equation of $y = f(x)$

$$y = \frac{1}{2}((-x)+2)^2 + 1 \Rightarrow y = \frac{1}{2}(-x+2)^2 + 1 \Rightarrow y = \frac{1}{2}[-1(x-2)]^2 + 1$$

Then factor out a "-1" from inside the brackets

Replace "x" with "-x" for horiz. reflection

$$\Rightarrow y = \frac{1}{2}(-1)^2(x-2)^2 + 1 \Rightarrow y = \frac{1}{2}(x-2)^2 + 1$$


Same resulting equation as replacing "x" with "x-4" !

(c) For graph of $y = h(x)$, (inverse) switch all points $(x, y) \rightarrow (y, x)$

$$(-4, 3) \rightarrow (3, -4)$$

$$(-2, 1) \rightarrow (1, -2)$$

$$(0, 3) \rightarrow (3, 0)$$



For equation of $y = h(x)$, switch x and y in the equation and isolate y .

$$y = \frac{1}{2}(x+2)^2 + 1$$

$$x = \frac{1}{2}(y+2)^2 + 1$$

$$x-1 = \frac{1}{2}(y+2)^2$$

$$2(x-1) = (y+2)^2$$

$$\begin{aligned} \text{sq. root both sides} \\ \sqrt{2(x-1)} &= \sqrt{(y+2)^2} \\ y+2 &= \sqrt{\pm 2(x-1)} \\ y &= \pm \sqrt{2(x-1)} - 2 \end{aligned}$$

$$h(x) = \pm \sqrt{2(x-1)} - 2$$

NOTE: without the "±", graph would only be a (sideways) half-parabola

WR Question 2

- (a) ① Vert. str. by factor of $\frac{3}{4}$, then vert. translation 6 units down

Next, vert shift 6 down....

$$y = \frac{3}{4}(2x^2 + 4x + 8) \rightarrow y = \frac{6}{4}x^2 + \frac{12}{4}x + \frac{24}{4} \rightarrow y = \frac{3}{2}x^2 + 3x + 6 \rightarrow y = \frac{3}{2}x^2 + 3x + 6 - 6 \rightarrow g(x) = \frac{3}{2}x^2 + 3x$$

- ② On $f(x)$ vertex is $(4, 2)$, then on $g(x)$ its $(1, 2)$ \rightarrow **horiz. shift 3 left**

– sign is outside for **vertical reflection**, which would make the y -coord of the vertex -2 . However, on $g(x)$ the y -coord is 2. So graph must also have been vertically translated 4 units up.

$$\rightarrow (x, y) \rightarrow (x - 3, -y + 4)$$

- ③ From mapping rule $(x, y) \rightarrow (2x, y)$ we can see that a horiz str, factor of $\frac{1}{2}$ was applied



Check on your calc!

So, working backward from $g(x)$ to $f(x)$, we should apply a horiz str, **factor of 2**.

$$f(x) = \sqrt{3(2x - 4)} \quad \text{Replace } x \text{ with } 2x \text{ in } g(x)$$

$$f(x) = \sqrt{3 * 2(x - 2)} \quad \text{Factor out the } 2 \text{ "inside"}$$

$$\rightarrow f(x) = \sqrt{6(x - 2)} \quad \text{or, alternatively } f(x) = \sqrt{6x - 12}$$



Horiz. str, factor of 2 (works!)




- (b) **RANGE of $f(x)$ is $\{y \in \mathbb{R}\}$** since $f(x)$ is a third degree (odd) polynomial function.... can also be determined by graphing

x -intercepts of $f(x)$ are $x = -3, x = 1, \text{ and } x = 2$

Can be determined from **factors** or by graphing

$$f(x) = 2(x + 3)(x - 1)(x - 2)$$

\downarrow \downarrow \downarrow
 $x = -3$ $x = 1$ $x = 2$



This is all for $f(x)$, the original function!

y -intercept of $f(x)$ is $y = 12$

Set $x = 0 \dots f(0) = 2(0 + 3)(0 - 1)(0 - 2)$

NOW, mapping rule here is $(x, y) \rightarrow (-x, 3y)$ (horiz refl, vert. str. of 3)

So.... x -intercepts of $g(x)$ are $x = -2, x = -1, \text{ and } x = 3$

Each becomes negative of original x -int.

The y -intercept of $g(x)$ is $y = 9$

Mult. y -int. of $f(x)$, which is 12, by $3/4$

The **RANGE** of $g(x)$ is $\{y \in \mathbb{R}\}$

No change to **RANGE** of $f(x)$, since it's all reals!

NOTE: We need not find an equation for $g(x)$ (though you may to help justify your answers)